



Language: Macedonian

Day: 1

Вторник, 12 Април, 2016

Задача 1. Нека n е непарен позитивен цел број, и нека x_1, x_2, \dots, x_n се ненегативни реални броеви. Докажи дека

$$\min_{i=1,2,\dots,n} (x_i^2 + x_{i+1}^2) \leq \max_{j=1,2,\dots,n} (2x_j x_{j+1})$$

каде што $x_{n+1} = x_1$.

Задача 2. Нека $ABCD$ е тетивен четириаголник, и нека неговите дијагонали AC и BD се сечат во точката X . Нека C_1, D_1 и M се средини на отсечките CX, DX и CD , соодветно. Правите AD_1 и BC_1 се сечат во точката Y , а правата MY ги сече дијагоналите AC и BD во различни точки E и F , соодветно. Докажи дека правата XY е тангента на опишаната кружница околу триаголникот EFX .

Задача 3. Нека m е позитивен цел број. Разгледуваме $4m \times 4m$ квадратна шема составена од единечни квадрати. За два различни единечни квадрати велеме дека се *сродни* ако тие се или во иста редица или се во иста колона од квадратната шема. Било кој единечен квадрат од квадратната шема не е сроден сам со себе. Некои единечни квадрати од квадратната шема се обоени во плаво, така што секој единечен квадрат од квадратната шема е сроден со најмалку два плави единечни квадрати. Да се определи минималниот можен број на плави единечни квадрати.

Language: Macedonian

Време за работа: 4 часа и 30 минути
Секоја задача се вреднува со 7 поени



Language: Macedonian

Day: 2

Среда, 13 Април, 2016

Задача 4. Кружниците ω_1 и ω_2 имаат еднакви радиуси и се сечат во две различни точки X_1 и X_2 . Кружницата ω е тангентна кон двете дадени кружници, при што кружницата ω_1 ја допира однадвор во точката T_1 , а кружницата ω_2 ја допира однатре во точката T_2 . Докажи дека правите X_1T_1 и X_2T_2 се сечат во точка која припаѓа на кружницата ω .

Задача 5. Нека k и n се цели броеви такви што $k \geq 2$ и $k \leq n \leq 2k - 1$. На шаховска табла со димензии $n \times n$ поставуваме правоаголници со димензии $1 \times k$ и $k \times 1$, така што секоја плочка препокрива точно k полиња од шаховската табла, и две поставени плочки не се преклопуваат. Плочки поставуваме се додека поставување на нова плочка не е можно. За секои k и n да се определи минималниот број на правоаголници кои може да се постават на погоре опишаниот начин.

Задача 6. Нека S е множество од сите позитивни цели броеви n такви што n^4 е делив со некој од броевите $n^2 + 1, n^2 + 2, \dots, n^2 + 2n$. Докажи дека во множеството S има бесконечно многу броеви од секој од облиците $7m, 7m + 1, 7m + 2, 7m + 5, 7m + 6$, а во S нема броеви од облик $7m + 3$ и $7m + 4$, каде m е цел број.

Language: Macedonian

Време за работа: 4 часа и 30 минути
Секоја задача се вреднува со 7 поени

EGMO 2016, Day 1 – Solutions

Problem 1. Let n be an odd positive integer, and let x_1, x_2, \dots, x_n be non-negative real numbers. Show that

$$\min_{i=1, \dots, n} (x_i^2 + x_{i+1}^2) \leq \max_{j=1, \dots, n} (2x_j x_{j+1}),$$

where $x_{n+1} = x_1$.

Solution. In what follows, indices are reduced modulo n . Consider the n differences $x_{k+1} - x_k$, $k = 1, \dots, n$. Since n is odd, there exists an index j such that $(x_{j+1} - x_j)(x_{j+2} - x_{j+1}) \geq 0$. Without loss of generality, we may and will assume both factors non-negative, so $x_j \leq x_{j+1} \leq x_{j+2}$. Consequently,


$$\min_{k=1, \dots, n} (x_k^2 + x_{k+1}^2) \leq x_j^2 + x_{j+1}^2 \leq 2x_{j+1}^2 \leq 2x_{j+1}x_{j+2} \leq \max_{k=1, \dots, n} (2x_k x_{k+1}).$$

Remark. If $n \geq 3$ is odd, and one of the x_k is negative, then the conclusion may no longer hold. This is the case if, for instance, $x_1 = -b$, and $x_{2k} = a$, $x_{2k+1} = b$, $k = 1, \dots, (n-1)/2$, where $0 \leq a < b$, so the string of numbers is

$$-b, b, a, b, a, \dots, b, a.$$

If n is even, the conclusion may again no longer hold, as shown by any string of alternate real numbers: a, b, a, b, \dots, a, b , where $a \neq b$.

Problem 2. Let $ABCD$ be a cyclic quadrilateral, and let diagonals AC and BD intersect at X . Let C_1 , D_1 and M be the midpoints of segments CX , DX and CD , respectively. Lines AD_1 and BC_1 intersect at Y , and line MY intersects diagonals AC and BD at different points E and F , respectively. Prove that line XY is tangent to the circle through E , F and X .



Solution. We are to prove that $\angle EXY = \angle EFX$; alternatively, but equivalently, $\angle AYX + \angle XAY = \angle BYF + \angle XBY$.

Since the quadrangle $ABCD$ is cyclic, the triangles XAD and XBC are similar, and since AD_1 and BC_1 are corresponding medians in these triangles, it follows that $\angle XAY = \angle XAD_1 = \angle XBC_1 = \angle XBY$.

Finally, $\angle AYX = \angle BYF$, since X and M are corresponding points in the similar triangles ABY and C_1D_1Y : indeed, $\angle XAB = \angle XDC = \angle MC_1D_1$, and $\angle XBA = \angle XCD = \angle MD_1C_1$.

Problem 3. Let m be a positive integer. Consider a $4m \times 4m$ array of square unit cells. Two different cells are *related* to each other if they are in either the same row or in the same column. No cell is related to itself. Some cells are coloured blue, such that every cell is related to at least two blue cells. Determine the minimum number of blue cells.

Solution 1 (Israel). The required minimum is $6m$ and is achieved by a diagonal string of m 4×4 blocks of the form below (bullets mark centres of blue cells):



In particular, this configuration shows that the required minimum does not exceed $6m$.

We now show that any configuration of blue cells satisfying the condition in the statement has cardinality at least $6m$.

Fix such a configuration and let m_1^r be the number of blue cells in rows containing exactly one such, let m_2^r be the number of blue cells in rows containing exactly two such, and let m_3^r be the number of blue cells in rows containing at least three such; the numbers m_1^c , m_2^c and m_3^c are defined similarly.

Begin by noticing that $m_3^c \geq m_1^r$ and, similarly, $m_3^r \geq m_1^c$. Indeed, if a blue cell is alone in its row, respectively column, then there are at least two other blue cells in its column, respectively row, and the claim follows.

Suppose now, if possible, the total number of blue cells is less than $6m$. We will show that $m_1^r > m_3^c$ and $m_1^c > m_3^r$, and reach a contradiction by the preceding: $m_1^r > m_3^c \geq m_1^c > m_3^r \geq m_1^r$.

We prove the first inequality; the other one is dealt with similarly. To this end, notice that there are no empty rows — otherwise, each column would contain at least two blue cells, whence a total of at least $8m > 6m$ blue cells, which is a contradiction. Next, count rows to get $m_1^r + m_2^r/2 + m_3^r/3 \geq 4m$,


and count blue cells to get $m_1^r + m_2^r + m_3^r < 6m$. Subtraction of the latter from the former multiplied by $3/2$ yields $m_1^r - m_3^r > m_2^r/2 \geq 0$, and the conclusion follows.

Solution 2. To prove that a minimal configuration of blue cells satisfying the condition in the statement has cardinality at least $6m$, consider a bipartite graph whose vertex parts are the rows and the columns of the array, respectively, a row and a column being joined by an edge if and only if the two cross at a blue cell. Clearly, the number of blue cells is equal to the number of edges of this graph, and the relationship condition in the statement reads: for every row r and every column c , $\deg r + \deg c - \epsilon(r, c) \geq 2$, where $\epsilon(r, c) = 2$ if r and c are joined by an edge, and $\epsilon(r, c) = 0$ otherwise.

Notice that there are no empty rows/columns, so the graph has no isolated vertices. By the preceding, the cardinality of every connected component of the graph is at least 4, so there are at most $2 \cdot 4m/4 = 2m$ such and, consequently, the graph has at least $8m - 2m = 6m$ edges. This completes the proof.

Remarks. The argument in the first solution shows that equality to $6m$ is possible only if $m_1^r = m_3^r = m_1^c = m_3^c = 3m$, $m_2^r = m_2^c = 0$, and there are no rows, respectively columns, containing four blue cells or more.

Consider the same problem for an $n \times n$ array. The argument in the second solution shows that the corresponding minimum is $3n/2$ if n is divisible by 4, and $3n/2 + 1/2$ if n is odd; if $n \equiv 2 \pmod{4}$, the minimum in question is $3n/2 + 1$. To describe corresponding minimal configurations C_n , refer to the minimal configurations C_2, C_3, C_4, C_5 below:



The case $n \equiv 0 \pmod{4}$ was dealt with above: a C_n consists of a diagonal string of $n/4$ blocks C_4 . If $n \equiv r \pmod{4}$, $r = 2, 3$, a C_n consists of a diagonal string of $\lfloor n/4 \rfloor$ blocks C_4 followed by a C_r , and if $n \equiv 1 \pmod{4}$, a C_n consists of a diagonal string of $\lfloor n/4 \rfloor - 1$ blocks C_4 followed by a C_5 .

Minimal configurations are not necessarily unique (two configurations being equivalent if one is obtained from the other by permuting the rows and/or the columns). For instance, if $n = 6$, the configurations below are both minimal:




EGMO 2016, Day 2 – Solutions

Problem 4. Two circles, ω_1 and ω_2 , of equal radius intersect at different points X_1 and X_2 . Consider a circle ω externally tangent to ω_1 at a point T_1 , and internally tangent to ω_2 at a point T_2 . Prove that lines X_1T_1 and X_2T_2 intersect at a point lying on ω .

Solution 1. Let the line X_kT_k and ω meet again at X'_k , $k = 1, 2$, and notice that the tangent t_k to ω_k at X_k and the tangent t'_k to ω at X'_k are parallel. Since the ω_k have equal radii, the t_k are parallel, so the t'_k are parallel, and consequently the points X'_1 and X'_2 coincide (they are not antipodal, since they both lie on the same side of the line T_1T_2). The conclusion follows.

Solution 2. The circle ω is the image of ω_k under a homothety h_k centred at T_k , $k = 1, 2$. The tangent to ω at $X'_k = h_k(X_k)$ is therefore parallel to the tangent t_k to ω_k at X_k . Since the ω_k have equal radii, the t_k are parallel, so $X'_1 = X'_2$; and since the points X_k , T_k and X'_k are collinear, the conclusion follows.



Solution 3. Invert from X_1 and use an asterisk to denote images under this inversion. Notice that ω_k^* is the tangent from X_2^* to ω^* at T_k^* , and the pole X_1 lies on the bisectrix of the angle formed by the ω_k^* , not containing ω^* . Letting $X_1T_1^*$ and ω^* meet again at Y , standard angle chase shows that Y lies on the circle $X_1X_2^*T_2^*$, and the conclusion follows.

Remarks. The product h_1h_2 of the two homotheties in the first solution is reflexion across the midpoint of the segment X_1X_2 , which lies on the line T_1T_2 .

Various arguments, involving similarities, radical axes, and the like, work equally well to prove the required result.

Problem 5. Let k and n be integers such that $k \geq 2$ and $k \leq n \leq 2k - 1$. Place rectangular tiles, each of size $1 \times k$ or $k \times 1$, on an $n \times n$ chessboard so that each tile covers exactly k cells, and no two tiles overlap. Do this until

no further tile can be placed in this way. For each such k and n , determine the minimum number of tiles that such an arrangement may contain.

Solution. The required minimum is n if $n = k$, and it is $\min(n, 2n - 2k + 2)$ if $k < n < 2k$.

The case $n = k$ being clear, assume henceforth $k < n < 2k$. Begin by describing maximal arrangements on the board $[0, n] \times [0, n]$, having the above mentioned cardinalities.

If $k < n < 2k - 1$, then $\min(n, 2n - 2k + 2) = 2n - 2k + 2$. To obtain a maximal arrangement of this cardinality, place four tiles, $[0, k] \times [0, 1]$, $[0, 1] \times [1, k + 1]$, $[1, k + 1] \times [k, k + 1]$ and $[k, k + 1] \times [0, k]$ in the square $[0, k] \times [0, k]$, stack $n - k - 1$ horizontal tiles in the rectangle $[1, k + 1] \times [k + 1, n]$, and erect $n - k - 1$ vertical tiles in the rectangle $[k + 1, n] \times [1, k + 1]$.

If $n = 2k - 1$, then $\min(n, 2n - 2k + 2) = n = 2k - 1$. A maximal arrangement of $2k - 1$ tiles is obtained by stacking $k - 1$ horizontal tiles in the rectangle $[0, k] \times [0, k - 1]$, another $k - 1$ horizontal tiles in the rectangle $[0, k] \times [k, 2k - 1]$, and adding the horizontal tile $[k - 1, 2k - 1] \times [k - 1, k]$.

The above examples show that the required minimum does not exceed the mentioned values.

To prove the reverse inequality, consider a maximal arrangement and let r , respectively c , be the number of rows, respectively columns, not containing a tile.

If $r = 0$ or $c = 0$, the arrangement clearly contains at least n tiles.

If r and c are both positive, we show that the arrangement contains at least $2n - 2k + 2$ tiles. To this end, we will prove that the rows, respectively columns, not containing a tile are consecutive. Assume this for the moment, to notice that these r rows and c columns cross to form an $r \times c$ rectangular array containing no tile at all, so $r < k$ and $c < k$ by maximality. Consequently, there are $n - r \geq n - k + 1$ rows containing at least one horizontal tile each, and $n - c \geq n - k + 1$ columns containing at least one vertical tile each, whence a total of at least $2n - 2k + 2$ tiles.

We now show that the rows not containing a tile are consecutive; columns are dealt with similarly. Consider a horizontal tile T . Since $n < 2k$, the nearest horizontal side of the board is at most $k - 1$ rows away from the row containing T . These rows, if any, cross the k columns T crosses to form a rectangular array no vertical tile fits in. Maximality forces each of these rows to contain a horizontal tile and the claim follows.

Consequently, the cardinality of every maximal arrangement is at least $\min(n, 2n - 2k + 2)$, and the conclusion follows.

Remarks. (1) If $k \geq 3$ and $n = 2k$, the minimum is $n + 1 = 2k + 1$

and is achieved, for instance, by the maximal arrangement consisting of the vertical tile $[0, 1] \times [1, k + 1]$ along with $k - 1$ horizontal tiles stacked in $[1, k + 1] \times [0, k - 1]$, another $k - 1$ horizontal tiles stacked in $[1, k + 1] \times [k + 1, 2k]$, and two horizontal tiles stacked in $[k, 2k] \times [k - 1, k + 1]$. This example shows that the corresponding minimum does not exceed $n + 1 < 2n - 2k + 2$. The argument in the solution also applies to the case $n = 2k$ to infer that for a maximal arrangement of minimal cardinality either $r = 0$ or $c = 0$, and the cardinality is at least n . Clearly, we may and will assume $r = 0$. Suppose, if possible, such an arrangement contains exactly n tiles. Then each row contains exactly one tile, and there are no vertical tiles. Since there is no room left for an additional tile, some tile T must cover a cell of the leftmost column, so it covers the k leftmost cells along its row, and there is then room for another tile along that row — a contradiction.

(2) For every pair (r, c) of integers in the range $2k - n, \dots, k - 1$, at least one of which is positive, say $c > 0$, there exists a maximal arrangement of cardinality $2n - r - c$.

Use again the board $[0, n] \times [0, n]$ to stack $k - r$ horizontal tiles in each of the rectangles $[0, k] \times [0, k - r]$ and $[k - c, 2k - c] \times [k, 2k - r]$, erect $k - c$ vertical tiles in each of the rectangles $[0, k - c] \times [k - r, 2k - r]$ and $[k, 2k - c] \times [0, k]$, then stack $n - 2k + r$ horizontal tiles in the rectangle $[k - c, 2k - c] \times [2k - r, n]$, and erect $n - 2k + c$ vertical tiles in the rectangle $[2k - c, n] \times [1, k + 1]$.

Problem 6. Let S be the set of all positive integers n such that n^4 has a divisor in the range $n^2 + 1, n^2 + 2, \dots, n^2 + 2n$. Prove that there are infinitely many elements of S of each of the forms $7m, 7m + 1, 7m + 2, 7m + 5, 7m + 6$ and no elements of S of the form $7m + 3$ or $7m + 4$, where m is an integer.

Solution. The conclusion is a consequence of the lemma below which actually provides a recursive description of S . The proof of the lemma is at the end of the solution.

Lemma. *The fourth power of a positive integer n has a divisor in the range $n^2 + 1, n^2 + 2, \dots, n^2 + 2n$ if and only if at least one of the numbers $2n^2 + 1$ and $12n^2 + 9$ is a perfect square.*

Consequently, a positive integer n is a member of S if and only if $m^2 - 2n^2 = 1$ or $m^2 - 12n^2 = 9$ for some positive integer m .

The former is a Pell equation whose solutions are $(m_1, n_1) = (3, 2)$ and

$$(m_{k+1}, n_{k+1}) = (3m_k + 4n_k, 2m_k + 3n_k), \quad k = 1, 2, 3, \dots$$

In what follows, all congruences are modulo 7. Iteration shows that $(m_{k+3}, n_{k+3}) \equiv (m_k, n_k)$. Since $(m_1, n_1) \equiv (3, 2)$, $(m_2, n_2) \equiv (3, -2)$, and

$(m_3, n_3) \equiv (1, 0)$, it follows that S contains infinitely many integers from each of the residue classes 0 and ± 2 modulo 7.

The other equation is easily transformed into a Pell equation, $m^2 - 12n^2 = 1$, by noticing that m and n are both divisible by 3, say $m = 3m'$ and $n = 3n'$. In this case, the solutions are $(m_1, n_1) = (21, 6)$ and

$$(m_{k+1}, n_{k+1}) = (7m_k + 24n_k, 2m_k + 7n_k), \quad k = 1, 2, 3, \dots$$

This time iteration shows that $(m_{k+4}, n_{k+4}) \equiv (m_k, n_k)$. Since $(m_1, n_1) \equiv (0, -1)$, $(m_2, n_2) \equiv (-3, 0)$, $(m_3, n_3) \equiv (0, 1)$, and $(m_4, n_4) \equiv (3, 0)$, it follows that S contains infinitely many integers from each of the residue classes 0 and ± 1 modulo 7.

Finally, since the n_k from the two sets of formulae exhaust S , by the preceding no integer in the residue classes ± 3 modulo 7 is a member of S .

We now turn to the proof of the lemma. Let n be a member of S , and let $d = n^2 + m$ be a divisor of n^4 in the range $n^2 + 1, n^2 + 2, \dots, n^2 + 2n$, so $1 \leq m \leq 2n$. Consideration of the square of $n^2 = d - m$ shows m^2 divisible by d , so m^2/d is a positive integer. Since $n^2 < d < (n+1)^2$, it follows that d is not a square; in particular, $m^2/d \neq 1$, so $m^2/d \geq 2$. On the other hand, $1 \leq m \leq 2n$, so $m^2/d = m^2/(n^2 + m) \leq 4n^2/(n^2 + 1) < 4$. Consequently, $m^2/d = 2$ or $m^2/d = 3$; that is, $m^2/(n^2 + m) = 2$ or $m^2/(n^2 + m) = 3$. In the former case, $2n^2 + 1 = (m-1)^2$, and in the latter, $12n^2 + 9 = (2m-3)^2$.

Conversely, if $2n^2 + 1 = m^2$ for some positive integer m , then $1 < m^2 < 4n^2$, so $1 < m < 2n$, and $n^4 = (n^2 + m + 1)(n^2 - m + 1)$, so the first factor is the desired divisor.

Similarly, if $12n^2 + 9 = m^2$ for some positive integer m , then m is odd, $n \geq 6$, and $n^4 = (n^2 + m/2 + 3/2)(n^2 - m/2 + 3/2)$, and again the first factor is the desired divisor.