

APMO 2021 Solution

1. Prove that for each real number $r > 2$, there are exactly two or three positive real numbers x satisfying the equation $x^2 = r[x]$.

Note: $[x]$ denotes the largest integer less than or equal to x

Solution 1 Let $r > 2$ be a real number. Let x be a positive real number such that $x^2 = r[x]$ with $[x] = k$. Since $x > 0$ and $x^2 = rk$, we also have $k > 0$. From $k \leq x < k + 1$, we get $k^2 \leq x^2 = rk < k^2 + 2k + 1 \leq k^2 + 3k$, hence $k \leq r < k + 3$, or $r - 3 < k \leq r$. There are at most three positive integers in the interval $(r - 3, r]$. Thus there are at most three possible values for k . Consequently, there are at most three positive solutions to the given equation.

Now suppose that k is a positive integer in the interval $[r - 2, r]$. There are at least two such positive integer. Observe that $k \leq \sqrt{rk} \leq \sqrt{(k + 2)k} < k + 1$ and so $rk = r[\sqrt{rk}]$. We conclude that the equation $x^2 = r[x]$ has at least two positive solutions, namely $x = \sqrt{rk}$ with $k \in [r - 2, r]$.

Marking Scheme

Solution 1

(+5 points) Proving that there are at most three positive solutions to the equation.

- (+1 point) Showing that $[x] \leq r$.
- (+3 point) Showing that $[x] \geq r - 3$.

(+2 points) Proving that there are at least two positive solutions to the equation.

- (+1 point) Showing that $x = \sqrt{r[r]}$ satisfy the equation.
- (+1 point) Showing that $x = \sqrt{r([r] - 1)}$ satisfy the equation.

Partial Point

(+1 point) Proving that there are exactly three positive solutions for each integer r .

2. For a polynomial P and a positive integer n , define P_n as the number of positive integer pairs (a, b) such that $a < b \leq n$ and $|P(a)| - |P(b)|$ is divisible by n .

Determine all polynomial P with integer coefficients such that for all positive integers n , $P_n \leq 2021$.

Solution There are two possible families of solutions:

- $P(x) = x + d$, for some integer $d \geq -2022$.
- $P(x) = -x + d$, for some integer $d \leq 2022$.

Suppose P satisfies the problem conditions. Clearly P cannot be a constant polynomial. Notice that a polynomial P satisfies the conditions if and only if $-P$ also satisfies them. Hence, we may assume the leading coefficient of P is positive. Then, there exists positive integer M such that $P(x) > 0$ for $x \geq M$.

Lemma 1. For any positive integer n , the integers $P(1), P(2), \dots, P(n)$ leave pairwise distinct remainders upon division by n .

Proof. Assume for contradiction that this is not the case. Then, for some $1 \leq y < z \leq n$, there exists $0 \leq r \leq n - 1$ such that $P(y) \equiv P(z) \equiv r \pmod{n}$. Since $P(an + b) \equiv P(b) \pmod{n}$ for all a, b integers, we have $P(an + y) \equiv P(an + z) \equiv r \pmod{n}$ for any integer a . Let A be a positive integer such that $An \geq M$, and let k be a positive integer such that $k > 2A + 2021$. Each of the $2(k - A)$ integers $P(An + y), P(An + z), P((A + 1)n + y), P((A + 1)n + z), \dots, P((k - 1)n + y), P((k - 1)n + z)$ leaves one of the k remainders

$$r, n + r, 2n + r, \dots, (k - 1)n + r$$

upon division by kn . This implies that at least $2(k - A) - k = k - 2A$ (possibly overlapping) pairs leave the same remainder upon division by kn . Since $k - 2A > 2021$ and all of the $2(k - A)$ integers are positive, we find more than 2021 pairs a, b with $a < b \leq kn$ for which $|P(b)| - |P(a)|$ is divisible by kn - hence, $P_{kn} > 2021$, a contradiction. □

Next, we show that P is linear. Assume that this is not the case, i.e., $\deg P \geq 2$. Then we can find a positive integer k such that $P(k) - P(1) \geq k$. This means that among the integers $P(1), P(2), \dots, P(P(k) - P(1))$, two of them, namely $P(k)$ and $P(1)$, leave the same remainder upon division by $P(k) - P(1)$ - contradicting the lemma (by taking $n = P(k) - P(1)$). Hence, P must be linear.

We can now write $P(x) = cx + d$ with $c > 0$. We prove that $c = 1$ by two ways.

Solution 1 If $c \geq 2$, then $P(1)$ and $P(2)$ leave the same remainder upon division by c , contradicting the Lemma. Hence $c = 1$. □

Solution 2 Suppose $c \geq 2$. Let n be a positive integer such that $n > 2cM$, $n \left(1 - \frac{3}{2c}\right) > 2022$ and $2c|n$. Notice that for any positive integers i such that $\frac{3n}{2c} + i < n$, $P\left(\frac{3n}{2c} + i\right) - P\left(\frac{n}{2c} + i\right) = n$. Hence, $\left(\frac{n}{2c} + i, \frac{3n}{2c} + i\right)$ satisfies the condition in the question for all positive integers i such that $\frac{3n}{2c} + i < n$. Hence, $P_n > 2021$, a contradiction. Then, $c = 1$. □

If $d \leq -2023$, then there are at least 2022 pairs $a < b$ such that $P(a) = P(b)$, namely $(a, b) = (1, -2d - 1), (2, -2d - 2), \dots, (-d - 1, -d + 1)$. This implies that $d \geq -2022$.

Finally, we verify that $P(x) = x + d$ satisfies the condition for any $d \geq -2022$. Fix a positive integer n . Note that $||P(b)| - |P(a)|| < n$ for all positive integers $a < b \leq n$, so the only pairs a, b for which $|P(b)| - |P(a)|$ could be divisible by n are those for which $|P(a)| = |P(b)|$. When $d \geq -2022$, there are indeed at most 2021 such pairs.

Marking Scheme

(+5 points) Proving all P that satisfy the solution must be linear.

- (+3 point) Proving the lemma.

(+2 points) Finding all linear solutions.

- (+1 point) Proving that if P is linear, P is of the form:
 - $P(x) = x + d$, for some integer $d \geq -2022$.
 - $P(x) = -x + d$, for some integer $d \leq 2022$.
- (+1 point) Verifying all linear solutions.

3. Let $ABCD$ be a cyclic convex quadrilateral and Γ be its circumcircle. Let E be the intersection of the diagonals AC and BD , let L be the center of the circle tangent to sides AB, BC , and CD , and let M be the midpoint of the arc BC of Γ not containing A and D . Prove that the excenter of triangle BCE opposite E lies on the line LM .

Solution 1

Let L be the intersection of the bisectors of $\angle ABC$ and $\angle BCD$. Let N be the E -excenter of $\triangle BCE$. Let $\angle BAC = \angle BDC = \alpha, \angle DBC = \beta$ and $\angle ACB = \gamma$.

We have the following:

$$\begin{aligned}\angle CBL &= \frac{1}{2}\angle ABC = 90^\circ - \frac{1}{2}\alpha - \frac{1}{2}\gamma \text{ and } \angle BCL = 90^\circ - \frac{1}{2}\alpha - \frac{1}{2}\beta, \\ \angle CBN &= 90^\circ - \frac{1}{2}\beta \text{ and } \angle BCN = 90^\circ - \frac{1}{2}\gamma, \\ \angle MBL &= \angle MBC + \angle CBL = 90^\circ - \frac{1}{2}\gamma \text{ and } \angle MCL = 90^\circ - \frac{1}{2}\beta, \\ \angle LCN &= \angle LBN = 180^\circ - \frac{1}{2}(\alpha + \beta + \gamma).\end{aligned}$$

Applying the sine rule to $\triangle MBL$ and $\triangle MCL$ we obtain

$$\frac{MB}{ML} = \frac{MC}{ML} = \frac{\sin \angle BLM}{\sin \angle MBL} = \frac{\sin \angle CLM}{\sin \angle MCL}.$$

It follows that

$$\frac{\sin \angle BLM}{\sin \angle CLM} = \frac{\sin \angle MBL}{\sin \angle MCL} = \frac{\cos(\gamma/2)}{\cos(\beta/2)}. \quad (1)$$

Now

$$\frac{\sin \angle BLM}{\sin \angle MLC} \cdot \frac{\sin \angle LCN}{\sin \angle NCB} \cdot \frac{\sin \angle NBC}{\sin \angle NBL} = \frac{\cos(\gamma/2)}{\cos(\beta/2)} \cdot \frac{\sin(90^\circ - \frac{1}{2}\beta)}{\sin(90^\circ - \frac{1}{2}\gamma)} = 1.$$

Hence LM, BN, CN are concurrent and therefore L, M, N are collinear.

Alternative proof

We proceed similarly as above until the equation (1).

We use the following lemma.

Lemma: If $\pi > \alpha, \beta, \gamma, \delta > 0, \alpha + \beta = \gamma + \delta < \pi$, and $\frac{\sin \alpha}{\sin \beta} = \frac{\sin \gamma}{\sin \delta}$, then $\alpha = \gamma$ and $\beta = \delta$.

Proof of Lemma: Let $\theta = \alpha + \beta = \gamma + \delta$. Then $\frac{\sin(\theta - \beta)}{\sin \beta} = \frac{\sin(\theta - \delta)}{\sin \delta}$.

$$\begin{aligned}\iff \sin(\theta - \beta) \sin \delta &= \sin(\theta - \delta) \sin \beta \\ \iff (\sin \theta \cos \beta - \sin \beta \cos \theta) \sin \delta &= (\sin \theta \cos \delta - \sin \delta \cos \theta) \sin \beta \\ \iff \sin \theta \cos \beta \sin \delta &= \sin \theta \cos \delta \sin \beta \\ \iff \sin \theta \sin(\beta - \delta) &= 0\end{aligned}$$

Since $0 < \theta < \pi$, then $\sin \theta \neq 0$. Therefore, $\sin(\beta - \delta) = 0$, and we must have $\beta = \delta$.

Applying the sine rule to $\triangle NBL$ and $\triangle NCL$ we obtain

$$\begin{aligned}\frac{NB}{NL} &= \frac{\sin \angle BLN}{\sin \angle LBN}, \\ \frac{NC}{NL} &= \frac{\sin \angle CLN}{\sin \angle LCN}.\end{aligned}$$

Since $\angle LBN = \angle LCN$, it follows that

$$\frac{\sin \angle BLN}{\sin \angle CLN} = \frac{NB}{NC} = \frac{\sin \angle BCN}{\sin \angle CBN} = \frac{\cos(\gamma/2)}{\cos(\beta/2)} = \frac{\sin \angle BLM}{\sin \angle CLM}.$$

By the lemma, it is concluded that $\angle BLM = \angle BLN$ and $\angle CLM = \angle CLN$. Therefore, L, M, N are collinear.

Solution 2

Denote by N the excenter of triangle BCE opposite E . Since BL bisects $\angle ABC$, we have $\angle CBL = \frac{\angle ABC}{2}$. Since M is the midpoint of arc BC , we have $\angle MBC = \frac{1}{2}(\angle MBC + \angle MCB)$ It follows by angle chasing that

$$\begin{aligned} \angle MBL &= \angle MBC + \angle CBL = \frac{1}{2}(\angle MBC + \angle MCB + \angle ABC) \\ &= \frac{1}{2}(\angle MBA + \angle MCB) = 90^\circ - \frac{\angle BCE}{2} = \angle BCN. \end{aligned}$$

Denote by X and Y the second intersections of lines BM and CM with the circumcircle of BCL , respectively. Since $\angle MBC = \angle MCB$, we have $BC \parallel XY$. It suffices to show that $BN \parallel XL$ and $CN \parallel YL$. Indeed, from this it follows that $\triangle BCN \sim \triangle XYL$, and therefore a homothety with center M that maps B to X and C to Y also maps N to L , implying that N lies on the line LM .

By symmetry, it suffices to show that $CN \parallel YL$, which is equivalent to showing that $\angle BCN = \angle XYL$. But we have $\angle BCN = \angle MBL = \angle XBL = \angle XYL$, completing the proof.

Marking Scheme

Solution 1

(+2 point) Showing the equalities: $\angle MBL = \angle BCN, \angle CBN = \angle MCL, \angle LBN = \angle LCN$.

(+5 points) Proving L, M, N are collinear.

- (+5 point) Proving that $\frac{\sin \angle BLM}{\sin \angle MLC} \cdot \frac{\sin \angle LCN}{\sin \angle NCB} \cdot \frac{\sin \angle NBC}{\sin \angle NBL} = 1$.

or

- (+1 point) Proving the lemma.
- (+3 point) Proving that $\frac{\sin \angle BLM}{\sin \angle MLC} = \frac{\sin \angle BLN}{\sin \angle NLC}$.
- (+1 point) Concluding that L, M, N are collinear from the previous equation.

(-1 point) For every minor important details

Solution 2

(+2 points) Showing the equalities: $\angle LBM = \angle BCN$ or $\angle CBN = \angle LCM$.

(+1 point) Constructing X, Y as described in the solution .

(+4 points) Proving that $\triangle XYL$ is homothetic with $\triangle BCN$.

(-1 point) For every minor important details.

Partial Solution: (+2 point) Claiming $\triangle XYL$ homothetic with $\triangle BCN$, or similar statement.


- Given a 32×32 table, we put a mouse (facing up) at the bottom left cell and a piece of cheese at several other cells. The mouse then starts moving. It moves forward except that when it reaches a piece of cheese, it eats a part of it, turns right, and continues moving forward. We say that a subset of cells containing cheese is good if, during this process, the mouse tastes each piece of cheese exactly once and then falls off the table. Show that:

- (a) No good subset consists of 888 cells.
- (b) There exists a good subset consisting of at least 666 cells.


Solution.

- (a) For the sake of contradiction, assume a good subset consisting of 888 cells exists. We call those *cheese-cells* and the other ones *gap-cells*. Observe that since each cheese-cell is visited once, each gap-cell is visited at most twice (once vertically and once horizontally). Define a finite sequence s whose i -th element is C if the i -th step of the mouse was onto a cheese-cell, and G if it was onto a gap-cell. By assumption, s contains 888 C 's. Note that s does not contain a contiguous block of 4 (or more) C 's. Hence s contains at least $888/3 = 296$ such C -blocks and thus at least 295 G 's. But since each gap-cell is traversed at most twice, this implies there are at least $\lceil 295/2 \rceil = 148$ gap-cells, for a total of $888 + 148 = 1036 > 32^2$ cells, a contradiction.
- (b) Let L_i, X_i be two $2^i \times 2^i$ tiles that allow the mouse to “turn left” and “cross”, respectively. In detail, the “turn left” tiles allow the mouse to enter at its bottom left cell facing up and to leave at its bottom left cell facing left. The “cross” tiles allow the mouse to enter at its top right facing down and leave at its bottom left facing left, while also to enter at its bottom left facing up and leave at its top right facing right.


(a) Basic tiles



(b) Inductive construction



(c) 16×16



Note that given two $2^i \times 2^i$ tiles L_i, X_i we can construct larger $2^{i+1} \times 2^{i+1}$ tiles L_{i+1}, X_{i+1} inductively as shown on in (b). The construction works because the path intersects itself (or the other path) only inside the smaller X -tiles where it works by induction.

For a tile T , let $|T|$ be the number of pieces of cheese in it. By straightforward induction, $|L_i| = |X_i| + 1$ and $|L_{i+1}| = 4 \cdot |L_i| - 1$. From the initial condition $|L_1| = 3$. We now easily compute $|L_2| = 11$, $|L_3| = 43$, $|L_4| = 171$, and $|L_5| = 683$. Hence we get the desired subset.

Another proof of (a).

Let X_N be the largest possible density of cheese-cells in a good subset on an $N \times N$ table. We will show that $X_N \leq 4/5 + o(1)$. Specifically, this gives $X_{32} \leq 817/1024$. We look at the (discrete analogue) of the winding number of the trajectory of the mouse. Since the mouse enters and leaves the table, for every 4 right turns in its trajectory there has to be a self-crossing. But each self-crossing requires a different empty square, hence $X_N \leq 4/5$.

Marking Scheme

(+2 points) Proving there is no good subset consists of 888 cells.

(+5 points) Proving there exists a good subset consisting of at least 666 cells.

- (+1 point) Constructing two 2×2 tiles that allow the mouse to “turn left” and “cross”.
- (+3 points) Showing inductive construction.
- (+1 point) Finishing calculation.

5. Determine all functions $f : \mathbb{Z} \rightarrow \mathbb{Z}$ such that $f(f(a) - b) + bf(2a)$ is a perfect square for all integers a and b .

Solution 1.

There are two families of functions which satisfy the condition:

- (1) $f(n) = \begin{cases} 0 & \text{if } n \text{ is even, and} \\ \text{any perfect square} & \text{if } n \text{ is odd} \end{cases}$
- (2) $f(n) = n^2$, for every integer n .

It is straightforward to verify that the two families of functions are indeed solutions. Now, suppose that f is any function which satisfies the condition that $f(f(a) - b) + bf(2a)$ is a perfect square for every pair (a, b) of integers. We denote this condition by $(*)$. We will show that f must belong to either Family (1) or Family (2).

Claim 1. $f(0) = 0$ and $f(n)$ is a perfect square for every integer n .

Proof. Plugging $(a, b) \rightarrow (0, f(0))$ in $(*)$ shows that $f(0)(f(0) + 1) = z^2$ for some integer z . Thus, $(2f(0) + 1 - 2z)(2f(0) + 1 + 2z) = 1$. Therefore, $f(0)$ is either -1 or 0 .

Suppose, for sake of contradiction, that $f(0) = -1$. For any integer a , plugging $(a, b) \rightarrow (a, f(a))$ implies that $f(a)f(2a) - 1$ is a square. Thus, for each $a \in \mathbb{Z}$, there exists $x \in \mathbb{Z}$ such that $f(a)f(2a) = x^2 + 1$. This implies that any prime divisor of $f(a)$ is either 2 or is congruent to $1 \pmod{4}$, and that $4 \nmid f(a)$, for every $a \in \mathbb{Z}$.

Plugging $(a, b) \rightarrow (0, 3)$ in $(*)$ shows that $f(-4) - 3$ is a square. Thus, there is $y \in \mathbb{Z}$ such that $f(-4) = y^2 + 3$. Since $4 \nmid f(-4)$, we note that $f(-4)$ is a positive integer congruent to $3 \pmod{4}$, but any prime dividing $f(-4)$ is either 2 or is congruent to $1 \pmod{4}$. This gives a contradiction. Therefore, $f(0)$ must be 0 .

For every integer n , plugging $(a, b) \rightarrow (0, -n)$ in $(*)$ shows that $f(n)$ is a square.

Replacing b with $f(a) - b$, we find that for all integers a and b ,

$$f(b) + (f(a) - b)f(2a) \text{ is a square.} \tag{**}$$

Now, let S be the set of all integers n such that $f(n) = 0$. We have two cases:

- Case 1: S is unbounded from above.

We claim that $f(2n) = 0$ for any integer n . Fix some integer n , and let $k \in S$ with $k > f(n)$. Then, plugging $(a, b) \mapsto (n, k)$ in $(**)$ gives us that $f(k) + (f(n) - k)f(2n) = (f(n) - k)f(2n)$ is a square. But $f(n) - k < 0$ and $f(2n)$ is a square by Claim 1. This is possible only if $f(2n) = 0$. In summary, $f(n) = 0$ whenever n is even and Claim 1 shows that $f(n)$ is a square whenever n is odd.

- Case 2: S is bounded from above.

Let T be the set of all integers n such that $f(n) = n^2$. We show that T is unbounded from above. In fact, we show that $\frac{p+1}{2} \in T$ for all primes p big enough.

Fix a prime number p big enough, and let $n = \frac{p+1}{2}$. Plugging $(a, b) \mapsto (n, 2n)$ in $(**)$ shows us that $f(2n)(f(n) - 2n + 1)$ is a square for any integer n . For p big enough, we have $2n \notin S$, so $f(2n)$ is a non-zero square. As a result, when p is big enough, $f(n)$ and $f(n) - 2n + 1 = f(n) - p$ are both squares. Writing $f(n) = k^2$ and $f(n) - p = m^2$ for some $k, m \geq 0$, we have

$$(k + m)(k - m) = k^2 - m^2 = p \implies k + m = p, k - m = 1 \implies k = n, m = n - 1.$$

Thus, $f(n) = k^2 = n^2$, giving us $n = \frac{p+1}{2} \in T$.

Next, for all $k \in T$ and $n \in \mathbb{Z}$, plugging $(a, b) \mapsto (n, k)$ in $(**)$ shows us that $k^2 + (f(n) - k)f(2n)$ is a square. But that means $(2k - f(2n))^2 - (f(2n))^2 - 4f(n)f(2n) = 4(k^2 + (f(n) - k)f(2n))$ is also a square. When k is large enough, we have $|f(2n)^2 - 4f(n)f(2n)| + 1 < |2k - f(2n)|$. As a result, we must have $f(2n)^2 = 4f(n)f(2n)$ and thus $f(2n) \in \{0, 4f(n)\}$ for all integers n .

Finally, we prove that $f(n) = n^2$ for all integers n . Fix n , and take $k \in T$ big enough such that $2k \notin S$. Then, we have $f(k) = k^2$ and $f(2k) = 4f(k) = 4k^2$. Plugging $(a, b) \mapsto (k, n)$ to $(**)$ shows us that $f(n) + (k^2 - n)4k^2 = (2k^2 - n)^2 + (f(n) - n^2)$ is a square. Since T is unbounded from above, we can take $k \in T$ such that $2k \notin S$ and also $|2k^2 - n| > |f(n) - n^2|$. This forces $f(n) = n^2$, giving us the second family of solution.

Another approach of Case 1.

Claim 2. One of the following is true.

- (i) For every integer n , $f(2n) = 0$.
- (ii) There exists an integer $K > 0$ such that for every integer $n \geq K$, $f(n) > 0$.

Proof. Suppose that there exists an integer $\alpha \neq 0$ such that $f(2\alpha) > 0$. We claim that for every integer $n \geq f(\alpha) + 1$, we have $f(n) > 0$.

For every $n \geq f(\alpha) + 1$, plugging $(a, b) \rightarrow (\alpha, f(\alpha) - n)$ in $(*)$ shows that $f(n) + (f(\alpha) - n)f(2\alpha)$ is a square, and in particular, is non-negative. Hence, $f(n) \geq (n - f(\alpha))f(2\alpha) > 0$, as desired.

If f belongs to Case (i), Claim 1 shows that f belongs to Family (1).

If f belongs to Case (ii), then S is bounded from above. From Case 2 we get $f(n) = n^2$.

Marking Scheme

(+1 point) Proving that $f(n)$ is a perfect square for every integer n .

(+2 point) Proving Case 1 or Claim 2.

(+4 points) Proving $f(n) = n^2$ when S is bounded from above.

- (+1 point) Proving $T = \{n : f(n) = n^2\}$ is unbounded.
- (+1 point) Proving $f(2n) = 4f(n)$.